

ON THE RATIONALITY OF MODULI SPACES OF POINTED CURVES

G. CASNATI AND C. FONTANARI

ABSTRACT. Motivated by several recent results on the geometry of the moduli spaces $\overline{\mathcal{M}}_{g,n}$ of stable curves of genus g with n marked points, we determine here their birational structure for small values of g and n by exploiting suitable plane models of the general curve. More precisely, we show that $\mathcal{M}_{g,n}$ is rational for $g = 2$ and $1 \leq n \leq 12$, $g = 3$ and $1 \leq n \leq 14$, $g = 4$ and $1 \leq n \leq 15$, $g = 5$ and $1 \leq n \leq 12$.

0. INTRODUCTION AND NOTATION

The geometry of algebraic curves varying in families is a very old and fascinating subject, which naturally leads to consider the moduli space $\mathcal{M}_g$ of smooth projective curves of genus g over the complex field $\mathbb{C}$.

In order to investigate $\mathcal{M}_g$ with the toolkit of algebraic geometry, the key ingredient is provided by its Deligne-Mumford compactification $\overline{\mathcal{M}}_g$ defined in terms of stable curves. Indeed, $\overline{\mathcal{M}}_g$ turns out to be a (mildly singular) projective variety and its boundary $\Delta = \overline{\mathcal{M}}_g \setminus \mathcal{M}_g$ has pure codimension one. Moreover, Δ presents a natural inductive structure, its irreducible components being obtained from moduli spaces of strictly smaller genus via the operation of gluing curves at pairs of points.

This remarkable fact suggests to consider more generally the moduli spaces $\mathcal{M}_{g,n}$ of smooth n -pointed curves of genus g , parametrizing curves together with an ordered n -tuple of distinct points, and their analogous compactifications $\overline{\mathcal{M}}_{g,n}$. Again, each irreducible component of the boundary $\overline{\mathcal{M}}_{g,n} \setminus \mathcal{M}_{g,n}$ is the image of either $\overline{\mathcal{M}}_{g-1,n+2}$ or $\overline{\mathcal{M}}_{g_1,n_1+1} \times \overline{\mathcal{M}}_{g_2,n_2+1}$ with $g_1 + g_2 = g$ and $n_1 + n_2 = n$ via a suitable gluing morphism. In order to exploit such an inductive pattern, the philosophy is to investigate these moduli spaces all together, even if one is mainly interested in $\overline{\mathcal{M}}_g$.

Indeed, the inductive approach has recently reached significant results both from a topological and a geometrical point of view: we cannot resist to mention at least the computation of the first few rational cohomology groups of $\overline{\mathcal{M}}_{g,n}$ by Arbarello and Cornalba in [A–C2] and the conjectural description of the ample cone of $\overline{\mathcal{M}}_{g,n}$ in purely combinatorial terms by Gibney, Keel, and Morrison in [G–K–M].

Such a context also motivates the effort of extending to the pointed case the many techniques and results available for $\overline{\mathcal{M}}_g$. For instance, the fact that its Kodaira dimension

2000 *Mathematics Subject Classification.* 14H10, 14H45, 14E08, 14L30.

Key words and phrases. Pointed curve, Moduli space, Rationality.

$\kappa(\overline{\mathcal{M}}_g)$ is non-negative for every $g \geq 23$ has been established in a long series of papers (see [H–M], [Hr], [E–H], [Fa1]). Along these lines, a nice contribution [Lg] by Logan determines for each g with $4 \leq g \leq 23$ a small integer $h(g)$ such that the Kodaira dimension $\kappa(\overline{\mathcal{M}}_{g,n})$ is non-negative for $n \geq h(g)$ (see also [Fa2] and [Fa3] for some recent improvements).

In the present paper instead we are going to prove the following

Main Theorem. *The moduli space $\mathcal{M}_{g,n}$ is rational for $g = 2$ and $1 \leq n \leq 12$, $g = 3$ and $1 \leq n \leq 14$, $g = 4$ and $1 \leq n \leq 15$, $g = 5$ and $1 \leq n \leq 12$.*

A few comments are in order here. The case $g = 0$ is trivial: indeed, $\mathcal{M}_{0,n}$ is birational to $(\mathbb{P}_{\mathbb{C}}^1)^{n-3}$ for every $n \geq 3$, hence it is rational whenever it is defined. The case $g = 1$ has been settled by P. Belorousski. In particular in [Be] it is proved via a nice geometric analysis of some plane cubic models of n -pointed elliptic curves that $\mathcal{M}_{1,n}$ is rational for $1 \leq n \leq 10$: for the sake of completeness we give a quick resumé of such a description in Section 1. We also point out that Belorousski's result is sharp: indeed, the fact that $H^{11,0}(\overline{\mathcal{M}}_{1,11}, \mathbb{C}) \neq 0$ (see for instance [G–P], Section 4.1), implies that $\mathcal{M}_{1,11}$ is not even unirational. More precisely, $\kappa(\overline{\mathcal{M}}_{1,11}) = 0$ and $\kappa(\overline{\mathcal{M}}_{1,n}) = 1$ for each $n \geq 12$ (see [B–F], Theorem 3).

In higher genus the state of the art is as follows. It is well-known that $\mathcal{M}_g$ is rational for $2 \leq g \leq 6$ (see respectively [Ig], [Ka4], [Ka2] and [SB1], [Ka3], [SB2]), and unirational for $7 \leq g \leq 14$ (see [A–S], [A–C1], [Se], [C–R], [Ve] and the references cited there).

The rationality of $\mathcal{M}_{g,1}$ is also known for $g = 2$ (see Section 6 of [Do]), $g = 3$ (see Section 10 of [Do], where the rationality of $\mathcal{M}_{3,1}$ is attributed to Shepherd-Barron without quoting any reference, and [Lo], Corollary 1.9, where an explicit proof is provided) and $g = 4$ (see Theorem B in [Ca]). Moreover, in Section 7 of [Lg] the author determines for $g = 2, \dots, 9, 11$ an integer $f(g)$ such that $\mathcal{M}_{g,n}$ is unirational for $n \leq f(g)$.

The Main Theorem above shows that unirationality can be improved to rationality at least for genus up to 5. We stress that for $g = 2$ we even cover the first case in which unirationality was not previously known and for $g \geq 3$ our bounds for n coincide exactly with the numbers $f(g)$ computed in [Lg], which turn out to be sharp at least in genus $g = 4$. We also point out that our approach relies on classical constructions of suitable plane models of the general curve of genus g , obtained from its canonical embedding by using the marked points. Hence we deduce the existence of a birational equivalence $\mathcal{M}_{g,n} \approx X/G$, where G is a linear group and X is a G -linearized projective bundle. Then we prove the rationality of $\mathcal{M}_{g,n}$ by checking the rationality of the quotient X/G via a case-by-case careful analysis exploiting several technical tools from rationality theory (for an overview of the standard methods see for instance [Do], [SB3], and Section 2.9 of [P–V]). Unfortunately, this natural strategy does not seem to apply for $g \geq 6$ and we leave the interested reader with the following

Main Question. *Is $\mathcal{M}_{g,n}$ rational for all $n \leq f(g)$, where $6 \leq g \leq 11$?*

The authors wish to thank Gilberto Bini and Roberto Notari for several stimulating conversations on the subject of the present paper.

Notation. We work over the field $\mathbb{C}$ of complex numbers. We denote by GL_k the general linear group of $k \times k$ matrices with entries in $\mathbb{C}$ and by PGL_k the projective linear group, i.e. GL_k modulo the subgroup of scalar matrices.

Let $\mathbb{C}[x_1, \dots, x_k]$ be the ring of polynomials in the variables $x_1, \dots, x_k$ with coefficients in $\mathbb{C}$, $\mathbb{C}[x_1, \dots, x_k]_d$ the vector space of degree d forms.

If $g_1, \dots, g_h$ are elements of a certain group (resp. vector space) G then $\langle g_1, \dots, g_h \rangle$ denotes the subgroup (resp. subspace) of G generated by $g_1, \dots, g_h$.

Projective r -space will be denoted by $\mathbb{P}_{\mathbb{C}}^r$: for each $i = 1, \dots, r+1$ we denote by E_i the point in $\mathbb{P}_{\mathbb{C}}^r$ all of whose coordinates but the i^{th} one are zero: $E_{r+2} \in \mathbb{P}_{\mathbb{C}}^r$ is the unit point, i.e. the point whose coordinates are all 1. If V is a vector space then $\mathbb{P}(V)$ is the associated projective space.

If X is a projective scheme, then a divisor D on X is always intended to be a Cartier divisor and $|D|$ denotes the complete linear system of divisors linearly equivalent to D .

A curve C is a projective scheme of dimension 1. If D is a divisor on C then $\varphi_{|D|}$ denotes the map induced on C by $|D|$. If $C, D \subset \mathbb{P}_{\mathbb{C}}^2$ are distinct plane curves, then $C \cdot D$ denotes their intersection cycle.

Let $\mathcal{M}_{g,n}$ be the coarse moduli space of smooth, projective n -pointed curves (briefly n -pointed curves in what follows) of genus g . A point of $\mathcal{M}_{g,n}$ is then an ordered $(n+1)$ -tuple $(C, p_1, \dots, p_n)$, where C is a smooth and connected curve of genus g and $p_1, \dots, p_n \in C$ are pairwise distinct point. As usual, we set $\mathcal{M}_g := \mathcal{M}_{g,0}$. A general curve of genus g is a curve corresponding to the general point of the irreducible scheme $\mathcal{M}_g$.

We denote isomorphisms by $\cong$ and birational equivalences by $\approx$.

For other definitions, results and notation we always refer to [Ht].

1. THE RATIONALITY OF $\mathcal{M}_{1,n}$ FOR $1 \leq n \leq 10$ (FOLLOWING BELOROUSSKI)

In this section we recall some known results about the rationality of $\mathcal{M}_{1,n}$ for $1 \leq n \leq 10$ due to P. Belorousski (see [Be]). In order to follow Belorousski's construction, fix four points A_1, A_2, A_3, A_4 in $\mathbb{P}_{\mathbb{C}}^2$ in linearly general position and three lines $r_{1,2}$, $r_{2,4}$ and $r_{3,4}$ such that the line $r_{i,j}$ passes through the points A_i and A_j .

1.1. The case $n \leq 2$. It is well-known that $\mathcal{M}_{1,1} \cong \mathbb{C}$ via the j -invariant, so the first non-trivial case is $n = 2$. In the space $\mathbb{C}[x_1, x_2, x_3]_3$ of cubic forms consider the subspace V_2 corresponding to cubics in $\mathbb{P}_{\mathbb{C}}^2$ passing through A_1, A_2, A_3, A_4 and tangent to $r_{1,2}$ at A_1 , to $r_{2,4}$ at A_2 , and to $r_{3,4}$ at A_4 . It is easy to see that $\mathbb{P}(V_2) \cong \mathbb{P}_{\mathbb{C}}^2$ and that the general member of $\mathbb{P}(V_2)$ is smooth. Hence there is a natural rational map $m_{1,2} : \mathbb{P}(V_2) \dashrightarrow \mathcal{M}_{1,2}$ taking a cubic with its points A_1 and A_2 to the corresponding 2-pointed curve of genus 1. As checked in [Be], Lemma 1.1.4, $m_{1,2}$ turns out to be birational, so $\mathcal{M}_{1,2}$ is rational.

1.2. The case $3 \leq n \leq 4$. Here the argument is very similar to the case $n = 2$. Define $\mathbb{P}(V_3) \subset \mathbb{P}_{\mathbb{C}}^9$ as the set of cubics in $\mathbb{P}_{\mathbb{C}}^2$ passing through A_1, A_2, A_3, A_4 and tangent to $r_{1,2}$ at A_1 and to $r_{2,4}$ at A_2 , and $\mathbb{P}(V_4) \subset \mathbb{P}_{\mathbb{C}}^9$ as the set of cubics in $\mathbb{P}_{\mathbb{C}}^2$ passing through A_1, A_2, A_3, A_4 and tangent to $r_{1,2}$ at A_1 . Exactly as in the previous subsection, there are natural isomorphisms $\mathbb{P}(V_3) \cong \mathbb{P}_{\mathbb{C}}^3$ and $\mathbb{P}(V_4) \cong \mathbb{P}_{\mathbb{C}}^4$ and rational maps $m_{1,3} : \mathbb{P}(V_3) \dashrightarrow \mathcal{M}_{1,3}$ and $m_{1,4} : \mathbb{P}(V_4) \dashrightarrow \mathcal{M}_{1,4}$ forgetting the plane embedding. Once again, in [Be] it is checked that both $m_{1,3}$ and $m_{1,4}$ are birational, hence the rationality of $\mathcal{M}_{1,3}$ and $\mathcal{M}_{1,4}$ follows.

1.3. The case $5 \leq n \leq 10$. Let V denote the vector space of cubic forms on $\mathbb{P}_{\mathbb{C}}^2$ vanishing at the points A_1, A_2, A_3, A_4 . On each general cubic curve $C \in V$ the line $r_{1,2}$ cuts out a

third point A_5 different from A_1 and A_2 , thus the choice of four points uniquely determines a fifth point on the elliptic curve. Consider now the incidence variety

$$X_n := \{ (f, A_6, \dots, A_n) \in \mathbb{P}(V) \times (\mathbb{P}_{\mathbb{C}}^2)^{n-5} \mid f(A_i) = 0, i = 6, \dots, n \}.$$

Since the dimension of V is 6, it follows that X_n is non-empty for $n \leq 10$. Moreover, Belorousski shows that X_n is irreducible and its general point corresponds to a smooth curve (see Lemma 2.5.1 below for a slightly weaker generalization). Hence there is a canonical rational map $m_{1,n} : X_n \dashrightarrow \mathcal{M}_{1,n}$ associating to a general point $(f, A_6, \dots, A_n) \in X_n$ the genus one curve $\{f = 0\}$ together with the n distinct points $(A_1, \dots, A_n)$. As proved by Belorousski, $m_{1,n}$ is indeed birational for all $n \geq 5$ (see [Be], Lemma 1.2.2) and X_n turns out to be rational for $5 \leq n \leq 10$ (see [Be], Lemma 1.2.3). Hence $\mathcal{M}_{1,n}$ is rational for all $n \leq 10$.

2. THE RATIONALITY OF $\mathcal{M}_{2,n}$ FOR $1 \leq n \leq 12$

Let C be a general curve of genus 2. Then C is hyperelliptic and it carries exactly one g_2^1 which is the canonical series. If we choose four, possibly coinciding, points $p_1, p_2, p_3, p_4 \in C$ then $D := p_1 + p_2 + p_3 + p_4$ defines a linear system $|D|$ which is a g_4^2 . By degree reasons $|D|$ has no base points, so we obtain a morphism $\varphi_{|D|} : C \rightarrow C_D \subseteq \mathbb{P}_{\mathbb{C}}^2$, whose image could be either an integral conic or a quartic divisor in $\mathbb{P}_{\mathbb{C}}^2$ with necessarily a double point by the genus formula for plane curves (see for instance [Ha], Example V.3.9.2). In the first case the g_4^2 should be composed with the unique canonical g_2^1 , which is absurd if the points in the support of D are general.

Remark 2.1. If $\Psi : C \xrightarrow{\sim} C'$ is an isomorphism of abstract curves of genus 2 such that $\Psi(p_i) = p'_i$, $i = 1, \dots, 4$ then Ψ carries $|D|$ to $|D'| := |p'_1 + p'_2 + p'_3 + p'_4|$, thus it induces a projectivity $\psi : \mathbb{P}_{\mathbb{C}}^2 \rightarrow \mathbb{P}_{\mathbb{C}}^2$ such that $\psi(C_D) = C'_{D'} := \varphi_{|D'|}(C')$ making the following diagram

$$(2.1.1) \quad \begin{array}{ccc} C & \xrightarrow{\Psi} & C' \\ \downarrow \varphi_{|D|} & & \downarrow \varphi_{|D'|} \\ C_D & \xrightarrow{\psi|_{C_D}} & C'_{D'} \end{array}$$

commutative. In particular $A_i := \varphi_{|D|}(p_i)$ is mapped by $\psi|_{C_D}$ to $A'_i := \varphi_{|D'|}(p'_i)$, $i = 1, \dots, 4$.

Conversely, each projectivity $\psi : \mathbb{P}_{\mathbb{C}}^2 \rightarrow \mathbb{P}_{\mathbb{C}}^2$ mapping C_D to $C'_{D'}$ and A_i to A'_i , $i = 1, \dots, 4$, is induced by an isomorphism $\Psi : C \xrightarrow{\sim} C'$ such that $\Psi(p_i) = p'_i$, $i = 1, \dots, 4$.

2.2. The case $n = 1$. If $(C, p_1) \in \mathcal{M}_{2,1}$ is general then let $D := 4p_1$. The morphism $\varphi_{|D|} : C \rightarrow \mathbb{P}_{\mathbb{C}}^2$ maps C to a quartic divisor $C_D \subseteq \mathbb{P}_{\mathbb{C}}^2$ with a 4-flex and by the genus formula C_D turns out to have also a double point. Up to projectivities we can assume that the singular point of C_D is $E_1 := [1, 0, 0]$, the 4-flex is $E_2 := [0, 1, 0]$ and the inflectional tangent line is $r := \{x_1 = 0\}$. Thus the equation of C_D is in

$$V := x_1^2 \mathbb{C}[x_2, x_3]_2 \oplus x_1 \mathbb{C}[x_2, x_3]_3 \oplus \mathbb{C}x_3^4.$$

Now the standard Bertini's theorem (see for instance [Ht], Theorem II.8.18 and Remark II.8.18.1) guarantees that the general element of V corresponds to an irreducible quartic divisor in $\mathbb{P}_{\mathbb{C}}^2$ with exactly one node at E_1 . Hence we have a rational dominant map $m_{2,1}: V \dashrightarrow \mathcal{M}_{2,1}$ and we are going to identify its fibres. Remark 2.1 implies that $f, f' \in V$ are in the same fibre of $m_{2,1}$ if and only if there is a projectivity fixing E_1, E_2 and r and sending $\{f = 0\}$ to $\{f' = 0\}$: thus it is represented by an element of the group

$$G := \left\{ \begin{pmatrix} \alpha_{1,1} & 0 & 0 \\ 0 & \alpha_{2,2} & \alpha_{2,3} \\ 0 & 0 & \alpha_{3,3} \end{pmatrix} \in \mathrm{GL}_3 \right\}.$$

It follows that $\mathcal{M}_{2,1} \approx V/G$.

2.3 The case $n = 2$. If $(C, p_1, p_2) \in \mathcal{M}_{2,2}$ is general then we can repeat the argument for the 1-pointed case with the divisor $D := 3p_1 + p_2$. We can thus assume that the singular point of C_D is E_1 and the flex is E_2 with inflectional tangent line $r := \{x_1 = 0\}$ cutting C_D also at $E_3 := [0, 0, 1]$. Thus the equation of C_D is in

$$V := x_1^2 \mathbb{C}[x_2, x_3]_2 \oplus x_1 \mathbb{C}[x_2, x_3]_3 \oplus \mathbb{C}x_2x_3^3.$$

Once again, we have a birational equivalence $\mathcal{M}_{2,2} \approx V/G$, where

$$G := \left\{ \begin{pmatrix} \alpha_{1,1} & 0 & 0 \\ 0 & \alpha_{2,2} & 0 \\ 0 & 0 & \alpha_{3,3} \end{pmatrix} \in \mathrm{GL}_3 \right\}.$$

2.4 The case $n = 3, 4$. If $(C, p_1, p_2, p_3) \in \mathcal{M}_{2,3}$ (resp. $(C, p_1, p_2, p_3, p_4) \in \mathcal{M}_{2,4}$) is general then we can repeat the argument for the 1-pointed case with the divisor $D := 2p_1 + p_2 + p_3$ (resp. $D := p_1 + p_2 + p_3 + p_4$). We can thus assume that the singular point of C_D is E_1 and that p_1, p_2, p_3 are respectively mapped to $E_2, E_3, E := [0, 1, 1]$. Thus the equation of C_D is in

$$V := x_1^2 \mathbb{C}[x_2, x_3]_2 \oplus x_1 \mathbb{C}[x_2, x_3]_3 \oplus \mathbb{C}x_2(x_2 - x_3)x_3^2$$

(resp. in

$$V := x_1^2 \mathbb{C}[x_2, x_3]_2 \oplus x_1 \mathbb{C}[x_2, x_3]_3 \oplus x_2x_3(x_2 - x_3)\mathbb{C}[x_2, x_3]_1).$$

Exactly as above, we obtain a birational equivalence $\mathcal{M}_{2,3} \approx V/G$ (resp. $\mathcal{M}_{2,4} \approx V/G$), where

$$G := \left\{ \begin{pmatrix} \alpha_{1,1} & 0 & 0 \\ 0 & \alpha_{2,2} & 0 \\ 0 & 0 & \alpha_{2,2} \end{pmatrix} \in \mathrm{GL}_3 \right\}.$$

2.5 The case $5 \leq n \leq 12$. If $(C, p_1, \dots, p_n) \in \mathcal{M}_{2,n}$ is general then we can repeat the argument for the 4-pointed case, so the equation of C_D is again in

$$V := x_1^2 \mathbb{C}[x_2, x_3]_2 \oplus x_1 \mathbb{C}[x_2, x_3]_3 \oplus x_2x_3(x_2 - x_3)\mathbb{C}[x_2, x_3]_1.$$

Consider the incidence variety

$$X := \{ (f, A_5, \dots, A_n) \in \mathbb{P}(V) \times (\mathbb{P}_{\mathbb{C}}^2)^{n-4} \mid f(A_i) = 0, i = 5, \dots, n \}.$$

Now we are going to define a rational dominant map $m_{2,n}: X \dashrightarrow \mathcal{M}_{2,n}$ and to investigate its fibres. To this purpose we need the following standard result, which will be helpful also in the next sections.

Lemma 2.5.1. *Let $V \subseteq \mathbb{C}[x_1, x_2, x_3]_d$ be a subspace of dimension v , fix $m \leq v - 1$ and define*

$$X' := \{ (f, A_1, \dots, A_m) \in \mathbb{P}(V) \times (\mathbb{P}_{\mathbb{C}}^2)^m \mid f(A_i) = 0, i = 1, \dots, m \}.$$

Then there exists a maximal non-empty open subset $\mathcal{U} \subseteq (\mathbb{P}_{\mathbb{C}}^2)^m$ such that $X := X' \cap (\mathbb{P}(V) \times \mathcal{U})$ is a subbundle of $\mathbb{P}(V) \times \mathcal{U}$ with fibre $\mathbb{P}_{\mathbb{C}}^{v-m-1}$ over $\mathcal{U}$, in particular it is irreducible. Moreover the projection $X \rightarrow \mathbb{P}(V)$ is dominant.

Proof. Let $\{f_1, \dots, f_v\}$ be a basis in V . Then each element $f \in V$ is of the form

$$f(x_1, x_2, x_3) = \sum_{i=1}^v \alpha_i f_i(x_1, x_2, x_3).$$

Let $x_{j,1}, x_{j,2}, x_{j,3}$ be coordinates on the j^{th} copy of $\mathbb{P}_{\mathbb{C}}^2$ inside $(\mathbb{P}_{\mathbb{C}}^2)^m$. Then $X' \subseteq \mathbb{P}(V) \times (\mathbb{P}_{\mathbb{C}}^2)^m$ is defined by the equations

$$\sum_{i=1}^v \alpha_i f_i(x_{j,1}, x_{j,2}, x_{j,3}) = 0, \quad j = 1, \dots, m.$$

In particular, the fibres over $(\mathbb{P}_{\mathbb{C}}^2)^m$ are projective spaces of dimension at least $v - m - 1$ linearly embedded inside $\mathbb{P}(V)$.

The dimension of the fibres over $(\mathbb{P}_{\mathbb{C}}^2)^m$ is an upper semicontinuous function and it is clear from the above equations that there are fibres of dimension exactly $v - m - 1$, hence there is a non-empty open subset $\mathcal{U} \subseteq (\mathbb{P}_{\mathbb{C}}^2)^m$ over which the fibre is $\mathbb{P}_{\mathbb{C}}^{v-m-1}$. By restriction of the tautological section of $\mathbb{P}(V) \times \mathcal{U}$, the subscheme $X := X' \cap (\mathbb{P}(V) \times \mathcal{U})$ is indeed a subbundle, so it is irreducible of dimension $\dim(\mathbb{P}_{\mathbb{C}}^2)^m + \dim \mathbb{P}_{\mathbb{C}}^{v-m-1} = v + m - 1$. Finally, since the fibres over a point of $X \rightarrow \mathbb{P}(V)$ are products of m times the corresponding curve, the projection must be dominant. $\square$

We now come back to our case. Here the general element in $\mathbb{P}(V)$ is irreducible and smooth outside its unique double point E_1 (by Bertini's theorem again). It follows that the same is true for the general point in $X \approx \mathbb{P}_{\mathbb{C}}^{12-n} \times (\mathbb{P}_{\mathbb{C}}^2)^{n-4}$, thus we obtain a rational map $m_{2,n}: X \dashrightarrow \mathcal{M}_{2,n}$. Let G be the image inside PGL_3 of

$$\left\{ \begin{pmatrix} \alpha_{1,1} & 0 & 0 \\ 0 & \alpha_{2,2} & 0 \\ 0 & 0 & \alpha_{2,2} \end{pmatrix} \in GL_3 \right\}.$$

Then the fibres of $m_{2,n}$ are the G -orbits of the points in X : in particular they have dimension 1. Since $\dim(X) = n + 4 = \dim(\mathcal{M}_{2,n}) + 1$ we conclude that $m_{2,n}$ is dominant and, as in the previous cases, we have $\mathcal{M}_{2,n} \approx X/G$.

We finally conclude this section with

Main Theorem for $g = 2$. *The moduli space $\mathcal{M}_{2,n}$ is rational for $1 \leq n \leq 12$.*

Proof. On one hand, if $1 \leq n \leq 4$ we have checked above that $\mathcal{M}_{2,n} \approx V/G$ where V is a linear representation of a solvable and connected algebraic group G . Thus the quotient on the right is rational by [Mi] or [Vi].

On the other hand, if $5 \leq n \leq 12$ we have checked above that $\mathcal{M}_{2,n} \approx X/G$ where X is a G -linearized projective bundle over the rational base $(\mathbb{P}_{\mathbb{C}}^2)^{n-4}$ with typical fibre $\mathbb{P}_{\mathbb{C}}^{12-n}$.

The scheme X is contained in the G -equivariant trivial projective bundle $\mathbb{P}(V) \times \mathcal{U}$. It is not difficult to check the existence of a G -invariant unisecant $L \subseteq \mathbb{P}(V) \times \mathcal{U}$ (i.e. a divisor intersecting the general fibre in a hyperplane). The scheme $L \cap X$ is then a G -invariant unisecant on X , hence X is G -equivariantly birationally equivalent to a G -equivariant vector bundle $\mathbb{C}^{12-n} \times (\mathbb{P}_{\mathbb{C}}^2)^{n-4}$ for $5 \leq n \leq 12$.

Obviously, the action of G on $(\mathbb{P}_{\mathbb{C}}^2)^{n-4}$ is almost free (i.e., the stabilizer in G of the general point of $(\mathbb{P}_{\mathbb{C}}^2)^{n-4}$ is trivial), thus $\mathcal{M}_{2,n} \approx X/G \approx \mathbb{C}^{12-n} \times (\mathbb{P}_{\mathbb{C}}^2)^{n-4}/G$ is a vector bundle over the base $(\mathbb{P}_{\mathbb{C}}^2)^{n-4}/G$ with $(12-n)$ -dimensional fibre by the results of Section 4 of [Do]. Moreover, G is a torus and acts linearly on $(\mathbb{P}_{\mathbb{C}}^2)^{n-4}$, thus $(\mathbb{P}_{\mathbb{C}}^2)^{n-4}/G$ is rational (e.g. since G is solvable and connected: see [Mi], [Vi]). $\square$

3. THE RATIONALITY OF $\mathcal{M}_{3,n}$ FOR $1 \leq n \leq 14$

Let C be a general curve of genus 3. Since C is not hyperelliptic, the canonical system $|K|$ is very ample, whence we obtain an embedding $\varphi_{|K|}: C \rightarrow C_K \subseteq \mathbb{P}_{\mathbb{C}}^2$. The image is a smooth quartic divisor.

Remark 3.1. If $\Psi: C \xrightarrow{\sim} C'$ is an isomorphism of abstract curves of genus 3 it carries the canonical system of C onto the one of C' . This is equivalent to saying that their canonical models C_K, C'_K are projectively equivalent.

3.2. The case $n = 1$. Let $(C, p_1) \in \mathcal{M}_{3,1}$. We can assume that $\varphi_K(p_1) = E_1$. Up to projectivities we can also assume that C_K is tangent to the line $r := \{x_3 = 0\}$ at the point E_1 . Thus the equation of C_K is in

$$V := \mathbb{C}x_1^3x_3 \oplus x_1^2\mathbb{C}[x_2, x_3]_2 \oplus x_1\mathbb{C}[x_2, x_3]_3 \oplus \mathbb{C}[x_2, x_3]_4.$$

As in Section 2.2 there is a rational dominant map $m_{3,1}: V \dashrightarrow \mathcal{M}_{3,1}$ and again Remark 3.1 allows us to identify its fibres with the orbits of the natural action of the group

$$G := \left\{ \begin{pmatrix} \alpha_{1,1} & \alpha_{1,2} & \alpha_{1,3} \\ 0 & \alpha_{2,2} & \alpha_{2,3} \\ 0 & 0 & \alpha_{3,3} \end{pmatrix} \in \mathrm{GL}_3 \right\}$$

on V . It follows that $\mathcal{M}_{3,1} \approx V/G$.

3.3 The case $n = 2, 3$. If $(C, p_1, p_2) \in \mathcal{M}_{3,2}$ (resp. $(C, p_1, p_2, p_3) \in \mathcal{M}_{3,3}$) we can assume that $\varphi_K(p_i) = E_i$, $i = 1, 2$ (resp. $i = 1, 2, 3$), up to projectivities. Thus the equation of C_K is in

$$V := x_1^3\mathbb{C}[x_2, x_3]_1 \oplus x_1^2\mathbb{C}[x_2, x_3]_2 \oplus \langle x_1, x_3 \rangle \mathbb{C}[x_2, x_3]_3$$

(resp. in

$$V := x_1^3 \mathbb{C}[x_2, x_3]_1 \oplus \langle x_1^2, x_2 x_3 \rangle \mathbb{C}[x_2, x_3]_2 \oplus x_1 \mathbb{C}[x_2, x_3]_3).$$

Once again, we have a birational equivalence $\mathcal{M}_{3,2} \approx V/G$ (resp. $\mathcal{M}_{3,3} \approx V/G$), where

$$G := \left\{ \begin{pmatrix} \alpha_{1,1} & 0 & \alpha_{1,3} \\ 0 & \alpha_{2,2} & \alpha_{2,3} \\ 0 & 0 & \alpha_{3,3} \end{pmatrix} \in \mathrm{GL}_3 \right\}$$

(resp.

$$G := \left\{ \begin{pmatrix} \alpha_{1,1} & 0 & 0 \\ 0 & \alpha_{2,2} & 0 \\ 0 & 0 & \alpha_{3,3} \end{pmatrix} \in \mathrm{GL}_3 \right\}).$$

3.4 The case $4 \leq n \leq 14$. If $(C, p_1, \dots, p_n) \in \mathcal{M}_{3,n}$ we can assume that $\varphi_K(p_i) = E_i$, $i = 1, 2, 3, 4$, so that the equation of C_D is in

$$V := \{ f \in x_1^3 \mathbb{C}[x_2, x_3]_1 \oplus \langle x_1^2, x_2 x_3 \rangle \mathbb{C}[x_2, x_3]_2 \oplus x_1 \mathbb{C}[x_2, x_3]_3 \mid f(1, 1, 1) = 0 \}.$$

Let us consider the incidence variety

$$X' := \{ (f, A_5, \dots, A_n) \in \mathbb{P}(V) \times (\mathbb{P}_{\mathbb{C}}^2)^{n-4} \mid f(A_i) = 0, i = 5, \dots, n \}$$

and consider the irreducible open subscheme $X \subseteq X'$ and the open set $\mathcal{U} \subseteq (\mathbb{P}_{\mathbb{C}}^2)^{n-4}$ defined in Lemma 2.5.1. Since the general element in $\mathbb{P}(V)$ is irreducible and smooth, the same is true for the general point in X . Since the unique automorphism of $\mathbb{P}_{\mathbb{C}}^2$ fixing the points E_i , $i = 1, 2, 3, 4$, is the identity we obtain that $\mathcal{M}_{3,n} \approx X \approx \mathbb{P}_{\mathbb{C}}^{14-n} \times (\mathbb{P}_{\mathbb{C}}^2)^{n-4}$ contained as a subbundle in $\mathbb{P}(V) \times (\mathbb{P}_{\mathbb{C}}^2)^{n-4}$.

Now the conclusion is analogous to the one of the previous section.

Main Theorem for $g = 3$. *The moduli space $\mathcal{M}_{3,n}$ is rational for $1 \leq n \leq 14$. $\square$*

4. THE RATIONALITY OF $\mathcal{M}_{4,n}$ FOR $1 \leq n \leq 15$

Let C be a general curve of genus 4. Then the canonical system $|K|$ is very ample, whence we obtain an embedding $\varphi_{|K|}: C \rightarrow C_K \subseteq \mathbb{P}_{\mathbb{C}}^3$. The image is a sextic curve which is the complete intersection of a smooth quadric Q with a cubic. Moreover two such sextic curves correspond to the same abstract curve if and only if they are projectively equivalent.

For each point $p_1 \in C$ the divisor $D := K - p_1$ defines a linear system $|D|$ which is a g_5^2 . The map $\varphi_{|D|}: C \rightarrow C_D \subseteq \mathbb{P}_{\mathbb{C}}^2$ coincides with the projection of C from p_1 . Its image C_D is a quintic divisor in $\mathbb{P}_{\mathbb{C}}^2$ passing through $A_1 := \varphi_{|D|}(p_1)$.

Proposition 4.1. *Under the above hypotheses, the curve C_D is an integral quintic with two nodes N_1, N_2 . The points N_1, N_2 and A_1 lie on a unique line r .*

Conversely, each integral quintic in $\mathbb{P}_{\mathbb{C}}^2$ with exactly two nodes N_1, N_2 is obtained in the above way.

Proof. We start by proving the second part of the statement. We recall that the blow up of the plane at two points is isomorphic to the blow up of Q at a point. Keeping this in mind, consider an integral quintic with two double points N_1, N_2 on the line r . Blowing up the plane at N_1, N_2 and blowing down the strict transform of r we obtain a smooth curve of genus 4 on Q passing through a fixed point. The genus formula for curves on a smooth quadric then shows that such a curve is the canonical model C_K of a suitable curve C of genus 4 with a fixed distinguished point p_1 and the restriction to C_K of the blow up/down map coincides with $\varphi_{|K-p_1|}$.

Conversely, C_D is obviously irreducible and it is also reduced by degree reasons. By the genus formula it must contain exactly two double points N_1, N_2 . If N_2 is infinitely near to N_1 , with a proper choice of the coordinates we can assume that $N_1 := [1, 0, 0]$ and that the tangent line at N_1 is $r := \{x_3 = 0\}$. Then the equation of C_D must lie in the space

$$\mathbb{C}x_1^3x_3^2 \oplus x_1^2x_3\mathbb{C}[x_2, x_3]_2 \oplus x_1\mathbb{C}[x_2, x_3]_4 \oplus \mathbb{C}[x_2, x_3]_5$$

and a parameter computation shows that the family of plane quintics with such a singular point is irreducible of dimension 17.

On the other hand, an analogous parameter computation shows that the family of plane quintics with two nodes is irreducible of dimension 18. Taking into account the first part of the proof, it follows that for a general curve C of genus 4 and for a general point $p_1 \in C$ the singularities of C_D are two distinct nodes.

Finally, if r is the line through N_1, N_2 , then it is easy to check that $r \cdot C_D = 2N_1 + 2N_2 + A_1$. $\square$

Remark 4.2. If $\Psi: C \xrightarrow{\sim} C'$ is an isomorphism of abstract curves of genus 4 mapping $p_1 \in C$ to $p'_1 \in C'$, then it carries the canonical system K of C onto the one K' of C' . Thus Ψ maps $|D| := |K - p_1|$ to $|D'| := |K' - p'_1|$. Hence there exists a projectivity $\psi: \mathbb{P}_{\mathbb{C}}^2 \rightarrow \mathbb{P}_{\mathbb{C}}^2$ such that $\psi(C_D) = C'_{D'} := \varphi_{|D'|}(C')$ and satisfying a commutative diagram of the form (2.1.1).

Conversely, each projectivity $\psi: \mathbb{P}_{\mathbb{C}}^2 \rightarrow \mathbb{P}_{\mathbb{C}}^2$ mapping C_D to $C'_{D'}$ and A_1 to A'_1 is induced by an isomorphism $\Psi: C \xrightarrow{\sim} C'$ such that $\Psi(p_1) = p'_1$.

The above Remark allows us to assume in what follows that the nodes are $N_i = E_i$, $i = 1, 2$, hence the line through the nodes is $r := \{x_3 = 0\}$. In particular, each projectivity $\psi: \mathbb{P}_{\mathbb{C}}^2 \rightarrow \mathbb{P}_{\mathbb{C}}^2$ mapping C_D to $C'_{D'}$ has to fix $\{E_1, E_2\}$.

4.3. The case $n = 1$. If $(C, p_1) \in \mathcal{M}_{4,1}$ and $D := K - p_1$, then the equation of C_D is in

$$U := x_1^3\mathbb{C}[x_2, x_3]_2 \oplus \langle x_1^2, x_1x_3, x_3^2 \rangle \mathbb{C}[x_2, x_3]_3.$$

Moreover, the line $r := \{x_3 = 0\}$ cuts out on C_D the divisor $2E_1 + 2E_2 + A_1$. By Bertini's theorem, there is a rational dominant map $m_{4,1}: U \dashrightarrow \mathcal{M}_{4,1}$ and Remark 4.2 allows us to identify its fibres with the orbits in U of the natural action of the group $H = H_0\langle\mu\rangle$, where

$$H_0 := \left\{ \begin{pmatrix} \alpha_{1,1} & 0 & \alpha_{1,3} \\ 0 & \alpha_{2,2} & \alpha_{2,3} \\ 0 & 0 & \alpha_{3,3} \end{pmatrix} \in \mathrm{GL}_3 \right\}, \quad \mu := \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

It follows that $\mathcal{M}_{4,1} \approx U/H$.

Let $G := T\langle\mu\rangle$, where T is the torus of diagonal matrices in GL_3 and

$$V := \langle x_1^3 x_2^2, x_1^3 x_3^2 \rangle \oplus x_1 x_2^2 \mathbb{C}[x_2, x_3]_2 \oplus \langle x_3^2, x_1^2 \rangle \mathbb{C}[x_2, x_3]_3.$$

We claim that V is a (H, G) -section of U in the sense of [Ka1] (see also Section 2.8 of [P-V]).

By definition, the first property to be checked is that V is G -dense in U . Indeed, let $f \in U$ be of the form

$$(4.3.1) \quad \begin{aligned} f(x_1, x_2, x_3) = & a_1 x_1^2 x_2^3 - 2a_2 x_1 x_2^3 x_3 + b_1 x_1^3 x_2^2 - 2b_2 x_1^3 x_2 x_3 + \\ & + \text{monomials not in } \langle x_1^2 x_2^3, x_1 x_2^3 x_3, x_1^3 x_2^2, x_1^3 x_2 x_3 \rangle. \end{aligned}$$

If we choose

$$h := \begin{pmatrix} 1 & 0 & a_2/a_1 \\ 0 & 1 & b_2/b_1 \\ 0 & 0 & 1 \end{pmatrix} \in H$$

then $h(f) \in V$ and we see that V is G -dense in U .

Next, we need to prove that if $f \in V$ is generic and $h \in H$ is such that $h(f) \in V$ then $h \in G$. Indeed, let $f \in V$ be as in formula (4.3.1): then $a_2 = b_2 = 0$ and if f is general enough we can suppose $a_1 b_1 \neq 0$. If

$$h := \begin{pmatrix} \alpha_{1,1} & 0 & \alpha_{1,3} \\ 0 & \alpha_{2,2} & \alpha_{2,3} \\ 0 & 0 & \alpha_{3,3} \end{pmatrix} \in H_0$$

then $h(f)$ contains the monomials $2a_1 \alpha_{1,1} \alpha_{1,2} \alpha_{2,2}^3 x_1 x_2^3 x_3 + 2b_1 \alpha_{1,1}^3 \alpha_{2,2} \alpha_{2,3} x_1^3 x_2 x_3$, hence it follows that $h(f) \in V$ if and only if $h \in T$. In a similar way one can argue for $h \in H_0 \mu$, so the claim is checked.

Now from the properties of sections (see Proposition 1.2 of [Ka1]) we deduce $\mathcal{M}_{4,1} \approx U/H \approx V/G$.

4.4 The case $n = 2$. If $(C, p_1, p_2) \in \mathcal{M}_{4,2}$, let again $D := K - p_1$ and consider $C_D \subseteq \mathbb{P}_{\mathbb{C}}^2$. We can assume that $\varphi_D(p_2) = E_3$. The equation of C_D is then in

$$V := \langle x_1^3, x_2 x_3^2 \rangle \mathbb{C}[x_2, x_3]_2 \oplus \langle x_1^2, x_1 x_3 \rangle \mathbb{C}[x_2, x_3]_3.$$

Once again, we have a birational equivalence $\mathcal{M}_{2,2} \approx V/G$, with $G = T\langle\mu\rangle$, T being as above the torus of diagonal matrices in GL_3 .

4.5 The case $3 \leq n \leq 15$. If $(C, p_1, \dots, p_n) \in \mathcal{M}_{4,n}$, then we can assume that $\varphi_D(p_i) = E_{i+1}$, $i = 2, 3$, so the equation of C_D is in

$$V := \{ f \in \langle x_1^3, x_2 x_3^2 \rangle \mathbb{C}[x_2, x_3]_2 \oplus \langle x_1^2, x_1 x_3 \rangle \mathbb{C}[x_2, x_3]_3 \mid f(1, 1, 1) = 0 \}.$$

Consider the irreducible scheme X and the non-empty open subset $\mathcal{U} \subseteq (\mathbb{P}_{\mathbb{C}}^2)^{n-3}$ defined in Lemma 2.5.1, so that $X \approx \mathbb{P}_{\mathbb{C}}^{15-n} \times \mathcal{U}$. The general element in $\mathbb{P}(V)$ is irreducible and smooth outside its double points E_1 and E_2 , hence the same is true for the general point in X . Since the only automorphisms of $\mathbb{P}_{\mathbb{C}}^2$ fixing the set $\{E_1, E_2\}$ and the points E_3, E_4 are the identity and μ , it follows that $\mathcal{M}_{4,n} \approx X/\langle\mu\rangle$.

The following theorem concludes this section.

Main Theorem for $g = 4$. *The moduli space $\mathcal{M}_{4,n}$ is rational for $1 \leq n \leq 15$.*

Proof. Consider the case $n = 1$ described in Section 4.3. We have thus a birational map $\mathcal{M}_{4,1} \approx V/G$. Now let $V_1 := \langle x_2^2 x_3^3, x_2 x_3^4, x_1^2 x_3^3, x_1 x_3^4 \rangle \subseteq V$. Since V_1 is G -invariant, there exists a decomposition $V = V_1 \oplus V_2$ for a suitable G -invariant subspace $V_2 \subseteq V$, in particular the projection $v: V \rightarrow V_1$ gives a vector bundle structure to V .

Lüroth's theorem guarantees that V_1/G is rational. We now compute the kernel K of the action of G on V_1 . Consider a general $c(x_1, x_2, x_3) := c_1 x_2^2 x_3^3 + c_2 x_2 x_3^4 + c_3 x_1^2 x_3^3 + c_4 x_1 x_3^4 \in V_1$. If

$$g := \begin{pmatrix} \alpha_{1,1} & 0 & 0 \\ 0 & \alpha_{2,2} & 0 \\ 0 & 0 & \alpha_{3,3} \end{pmatrix} \in T$$

is in the stabilizer G_c of c inside G , then $\alpha_{2,2}^2 \alpha_{3,3}^3 = \alpha_{2,2} \alpha_{3,3}^4 = \alpha_{1,1}^2 \alpha_{3,3}^3 = \alpha_{1,1} \alpha_{3,3}^4 = 1$, thus $\alpha_{1,1} = \alpha_{2,2} = \alpha_{3,3}$ and $\alpha_{1,1}^5 = 1$. A similar computation shows that no elements of the form

$$g := \begin{pmatrix} 0 & \alpha_{1,1} & 0 \\ \alpha_{2,2} & 0 & 0 \\ 0 & 0 & \alpha_{3,3} \end{pmatrix} \in T\mu$$

can be in the stabilizer of c . We conclude that K coincides with the normal subgroup of scalar fifth roots of the identity: in particular it coincides with the kernel of the action of G on V . Thus $\mathcal{M}_{4,1} \approx V/G$ is rational by the results of Section 4 in [Do].

Now consider the case $n = 2$ described in Section 4.4: once again, we have a birational map $\mathcal{M}_{4,2} \approx V/G$ and letting $V_1 := \langle x_2^2 x_3^3, x_2 x_3^4, x_1^2 x_3^3, x_1 x_3^4 \rangle \subseteq V$ we get a decomposition $V = V_1 \oplus V_2$ into G -invariant subspaces. Thus we can prove the statement exactly as in the previous case.

Finally consider the case $n \geq 3$ described in Section 4.5. In this case, $\mathcal{M}_{4,n} \approx X/\langle \mu \rangle$. The scheme $X \approx \mathbb{P}_{\mathbb{C}}^{15-n} \times \mathcal{U}$ is a subbundle of $\mathbb{P}(V) \times \mathcal{U}$ which obviously contains a $\langle \mu \rangle$ -invariant unisecant L (i.e. a divisor intersecting the general fibre in a hyperplane). Thus $L \cap X$ is still a $\langle \mu \rangle$ -invariant unisecant on X . Hence there exists a $\langle \mu \rangle$ -equivariant birational equivalence $X \approx \mathbb{C}^{15-n} \times (\mathbb{P}_{\mathbb{C}}^2)^{n-3}$ which is a $\langle \mu \rangle$ -equivariant vector bundle. The group $\langle \mu \rangle$ acts on $(\mathbb{P}_{\mathbb{C}}^2)^{n-3}$ by exchanging the first two coordinates in each factor, so its action is trivially almost free. Moreover, this description gives us an obvious $\langle \mu \rangle$ -equivariant birational equivalence $(\mathbb{P}_{\mathbb{C}}^2)^{n-3} \approx (\mathbb{C}^2)^{n-3}$. Once again, $\langle \mu \rangle$ exchanges the coordinates in each factor of $(\mathbb{C}^2)^{n-3}$, thus $(\mathbb{C}^2)^{n-3}$ is a linear representation of the abelian group $\langle \mu \rangle \cong \mathbb{Z}_2$ and the quotient is rational (the classical reference for this elementary fact is [Fi]). Now we conclude via the already quoted results in [Do]. $\square$

5. THE RATIONALITY OF $\mathcal{M}_{5,n}$ FOR $1 \leq n \leq 12$

Let C be a general curve of genus 5. Then the canonical system $|K|$ is very ample, whence we obtain an embedding $\varphi_{|K|}: C \rightarrow C_K \subseteq \mathbb{P}_{\mathbb{C}}^4$. The image is an octic curve which is the base locus of a net of quadrics.

For each pair of possibly coinciding general points $p_1, p_2 \in C$ the divisor $D := K - p_1 - p_2$ defines a linear system $|D|$ which is a g_6^2 . The map $\varphi_{|D|}: C \rightarrow C_D \subseteq \mathbb{P}_{\mathbb{C}}^2$ coincides with the

projection of C from the line joining p_1 and p_2 (if $p_1 = p_2$ it coincides with the projection from the tangent line to C at the point p_1). Its image C_D is a sextic divisor in $\mathbb{P}_{\mathbb{C}}^2$ passing through the points $A_i := \varphi_{|D|}(p_i)$.

Proposition 5.1. *Under the above hypotheses, the curve C_D is an integral sextic with five nodes $N_1, \dots, N_5$. The points $N_1, \dots, N_5$ and A_1, A_2 lie on a unique integral conic $\widehat{C}$. If $A_1 = A_2$ such a conic is tangent to C at A_1 .*

Conversely, each integral sextic in $\mathbb{P}_{\mathbb{C}}^2$ with exactly five nodes $N_1, \dots, N_5$ lying on an integral conic $\widehat{C}$ is obtained in the above way.

Proof. We start by proving the second part of the statement. Consider an integral sextic with five nodes $N_1, \dots, N_5$ in general position. Thus the blow up of the plane at the points $N_1, \dots, N_5$ is a quartic del Pezzo surface Σ . If we consider the anticanonical embedding of Σ inside $\mathbb{P}_{\mathbb{C}}^4$, then the strict transform $\widetilde{C}$ of the curve is smooth and it is cut out on Σ by a quadric. Since Σ is the complete intersection of two quadrics, $\widetilde{C}$ is the smooth complete intersection of three quadrics in $\mathbb{P}_{\mathbb{C}}^4$ i.e. it is the canonical model C_K of a suitable curve C of genus 5. The points $N_1, \dots, N_5$ lie on a unique integral conic $\widehat{C}$ cutting out only two other, possibly coinciding, points on our curve. The strict transform of $\widehat{C}$ is a line on Σ intersecting C_K at two, possibly coinciding, points p_1, p_2 . Then the restriction to C_K of the blow up map coincides with $\varphi_{|K-p_1-p_2|}$.

Conversely, it is clear that C_D is irreducible. In order to prove it is integral it then suffices to check it is reduced. If C_D is a triple conic (resp. a double smooth cubic, a double singular cubic) then C would be trigonal (resp. bielliptic, hyperelliptic), thus C would not be general.

Thus we can assume that C_D is actually integral. Clearly C_D is singular and by the genus formula it carries at most triple points as singularities. Since C , being general, is not trigonal, then C_D carries only double points, which turn out to be exactly five (possibly infinitely near), again by the genus formula. We denote them by $N_1, \dots, N_5$. Since C_D is integral such five nodes $N_1, \dots, N_5$ lie necessarily on a reduced conic $\widehat{C}$. If $\widehat{C}$ is integral then it is clearly unique. If $\widehat{C}$ is the union of two lines, one of them contains three of the five nodes $N_1, \dots, N_5$, the other one the remaining two points, hence it is unique also in this case.

Sextics with five assigned double points on a reduced conic $\widehat{C}$ form a linear system of dimension 21. If $\widehat{C}$ is reducible then the points $N_1, \dots, N_4$ can be chosen freely, the remaining one N_5 on the line joining N_1 and N_2 . Thus the locus of sextics with a reducible $\widehat{C}$ is dominated by a projective bundle with fibre $\mathbb{P}_{\mathbb{C}}^{12}$ over the product $(\mathbb{P}_{\mathbb{C}}^2)^4 \times \mathbb{P}_{\mathbb{C}}^1$, hence it is irreducible and its dimension is 21. Analogously, other standard parameter computations show that each family of such plane integral sextics with infinitely near double points has dimension at most 21.

Similarly one can also check that the locus of plane integral sextics with five pairwise distinct nodes on an integral conic is irreducible of dimension 22. Since we have proved above that each plane sextic with such a second configuration of singularities is the projection from a suitable secant line of its canonical model, it follows that for a general curve C of genus 5 and for general points $p_1, p_2 \in C$ the singularities of C_D are five nodes.

By construction, for each general line $\ell \subseteq \mathbb{P}_{\mathbb{C}}^2$ the divisor $\ell \cdot C_D + A_1 + A_2$ is in the

canonical system. Since the canonical system is cut out on C_D by the adjoints of degree 3 we have the linear equivalence on C_D

$$\ell \cdot C_D + A_1 + A_2 \sim \ell \cdot C_D + \widehat{C} \cdot C_D - 2 \sum_{i=1}^5 N_i.$$

Thus $A_1 + A_2 \sim \widehat{C} \cdot C_D - 2 \sum_{i=1}^5 N_i$ and since C_D is not hyperelliptic we have $\widehat{C} \cdot C_D = A_1 + A_2 + 2 \sum_{i=1}^5 N_i$, i.e. $A_1, A_2 \in \widehat{C}$. $\square$

Remark 5.2. As in the genus 4 case, if $\Psi: C \xrightarrow{\sim} C'$ is an isomorphism of abstract curves of genus 5 mapping $p_i \in C$ to $p'_i \in C'$, then it carries the canonical system K of C onto the one K' of C' . Thus Ψ maps $|D| := |K - p_1 - p_2|$ to $|D'| := |K' - p'_1 - p'_2|$. Hence there exists a projectivity $\psi: \mathbb{P}_{\mathbb{C}}^2 \rightarrow \mathbb{P}_{\mathbb{C}}^2$ sending C_D to $C'_{D'} := \varphi_{|D'|}(C')$ and inducing a commutative diagram of the form (2.1.1).

Conversely, each projectivity $\psi: \mathbb{P}_{\mathbb{C}}^2 \rightarrow \mathbb{P}_{\mathbb{C}}^2$ mapping C_D to $C'_{D'}$ and A_i to A'_i is induced by an isomorphism $\Psi: C \xrightarrow{\sim} C'$ such that $\Psi(p_i) = p'_i$, $i = 1, 2$.

The above remark allows us to assume in what follows that $A_1 = E_1$ (resp. $A_i = E_i$, $i = 1, 2$) and $\widehat{C} := \{x_3^2 - x_1 x_2 = 0\}$. Notice that $\widehat{C}$ is the image of the Veronese embedding $v: \mathbb{P}_{\mathbb{C}}^1 \rightarrow \mathbb{P}_{\mathbb{C}}^2$ given by $[s_1, s_2] \mapsto [s_1^2, s_2^2, s_1 s_2]$. In particular, two divisors $C_D, C'_{D'} \subseteq \mathbb{P}_{\mathbb{C}}^2$ turn out to be projectively equivalent if and only if they are mapped one onto the other by an automorphism of $\mathbb{P}_{\mathbb{C}}^2$ fixing A_1 (resp. A_i , $i = 1, 2$) and $\widehat{C}$. Every such automorphism is induced by an automorphism of $\widehat{C} \cong \mathbb{P}_{\mathbb{C}}^1$ fixing $v^{-1}(A_1)$ (resp. $v^{-1}(A_i)$, $i = 1, 2$), since its well known restriction gives an isomorphism between the group of automorphisms of $\mathbb{P}_{\mathbb{C}}^2$ fixing $\widehat{C}$ and the group of automorphisms of $\widehat{C} \cong \mathbb{P}_{\mathbb{C}}^1$.

The five nodes correspond to an unordered 5-tuple of pairwise distinct points $S_1, \dots, S_5$ in $\mathbb{P}_{\mathbb{C}}^1 \setminus \{E_1\}$ (resp. $\mathbb{P}_{\mathbb{C}}^1 \setminus \{E_1, E_2\}$). The set of such 5-tuples can be naturally identified with the open subset $\mathbb{P}_5^0 \subseteq \mathbb{P}_5 := \mathbb{P}(\mathbb{C}[s_1, s_2]_5)$ corresponding to polynomials without multiple roots and not divisible by s_2 (resp. either s_1 or s_2).

5.3. The case $n = 1$. If $(C, p_1) \in \mathcal{M}_{5,1}$ and $D := K - 2p_1$, then C_D is a sextic tangent to the line $\{x_2 = 0\}$ at the point E_1 , hence it is defined by a polynomial in

$$V := \mathbb{C}x_1^5 x_2 \oplus x_1^4 \mathbb{C}[x_2, x_3]_2 \oplus x_1^3 \mathbb{C}[x_2, x_3]_3 \oplus x_1^2 \mathbb{C}[x_2, x_3]_4 \oplus x_1 \mathbb{C}[x_2, x_3]_5 \oplus \mathbb{C}[x_2, x_3]_6.$$

If we consider the incidence variety

$$Y := \{ (f, S_1, \dots, S_5) \in \mathbb{P}(V) \times \mathbb{P}_5^0 \mid N_i := v(S_i) \text{ is a node of } \{f = 0\}, i = 1, \dots, 5 \}$$

with projection $\pi_Y: Y \rightarrow \mathbb{P}_5^0$, then the above description shows that the equation of C_D belongs necessarily to a fibre of π_Y . Conversely,

Lemma 5.3.1. *The scheme Y is a subbundle of $\mathbb{P}(V) \times \mathbb{P}_5^0$ with typical fibre $\mathbb{P}_{\mathbb{C}}^{10}$ over $\mathbb{P}_5^0$, in particular it is irreducible.*

Moreover the general point in Y corresponds to an irreducible sextic curve tangent to the line r at the point E_1 , containing exactly five nodes on $\widehat{C}$ and no other singularities.

Proof. The first part of the above statement can be checked by imitating the proof of Lemma 2.5.1: in this case, using reducible curves it is easy to see that the dimension of the fibres of π_Y over $\mathbb{P}_5^0$ is constant. The smoothness of the curve corresponding to the general point of Y can be easily proved by applying Bertini's theorem to suitable reducible curves singular at the points $N_1, \dots, N_5$. $\square$

Thus there is a rational dominant map $m_{5,1}: Y \dashrightarrow \mathcal{M}_{5,1}$: Remark 5.2 allows us to identify its fibres with the orbits in Y of the natural action of the stabilizer in GL_2 of the point $[1, 0] = v^{-1}(A_1)$, which is the image H inside PGL_2 of

$$\left\{ \begin{pmatrix} \alpha_{1,1} & \alpha_{1,2} \\ 0 & \alpha_{2,2} \end{pmatrix} \in \mathrm{GL}_2 \right\}.$$

It follows that $\mathcal{M}_{5,1} \approx Y/H$.

Let $\overline{\mathbb{P}}_5^0 := \mathbb{P}(\langle s_1^5, s_1^3 s_2^2, s_1^2 s_2^3, s_1 s_2^4, s_2^5 \rangle) \cap \mathbb{P}_5^0$, and consider the incidence variety $X := \pi_Y^{-1}(\overline{\mathbb{P}}_5^0)$ with projection $\pi := (\pi_Y)|_X: X \rightarrow \overline{\mathbb{P}}_5^0$. The variety X is generically a projective bundle with typical fibre $\mathbb{P}_{\mathbb{C}}^{10}$ over the rational base $B = \overline{\mathbb{P}}_5^0$.

Let $T \cong \mathbb{C}^* \subseteq \mathrm{PGL}_2$ be the image of the torus of diagonal matrices. We claim that X is a (H, T) -section of Y . Since $X = \pi_Y^{-1}(\overline{\mathbb{P}}_5^0)$ is irreducible and it dominates $\overline{\mathbb{P}}_5^0$, by Proposition 1.2 of [Ka1] it is sufficient to check that $\overline{\mathbb{P}}_5^0$ is a (H, T) -section of $\mathbb{P}_5^0$: this last assertion is an easy exercise using Tschirnhaus transformations.

We conclude that $\mathcal{M}_{5,1} \approx Y/H \approx X/T$.

5.4 The case $n = 2$. If $(C, p_1, p_2) \in \mathcal{M}_{5,2}$, set $D := K - p_1 - p_2$ and consider $C_D \subseteq \mathbb{P}_{\mathbb{C}}^2$. C_D is a sextic through the points E_1, E_2 , hence it is defined by a polynomial in

$$V := x_1^5 \mathbb{C}[x_2, x_3]_1 \oplus x_1^4 \mathbb{C}[x_2, x_3]_2 \oplus x_1^3 \mathbb{C}[x_2, x_3]_3 \oplus x_1^2 \mathbb{C}[x_2, x_3]_4 \oplus \langle x_1, x_3 \rangle \mathbb{C}[x_2, x_3]_5.$$

Once again, define the incidence variety

$$X := \{ (f, S_1, \dots, S_5) \in \mathbb{P}(V) \times \mathbb{P}_5^0 \mid N_i := v(S_i) \text{ is a node on } \{ f = 0 \}, i = 1, \dots, 5 \}.$$

As in the previous case, the following result holds.

Lemma 5.4.1. *The scheme X is a subbundle of $\mathbb{P}(V) \times \mathbb{P}_5^0$ with typical fibre $\mathbb{P}_{\mathbb{C}}^{10}$ over $\mathbb{P}_5^0$, in particular it is irreducible.*

Moreover the general point in Y corresponds to an irreducible sextic curve through the points E_1, E_2 , carrying exactly five nodes on $\widehat{C}$ and no other singularities. $\square$

As in Section 5.3, $\mathcal{M}_{5,2} \approx X/T$.

5.5 The case $3 \leq n \leq 12$. If $(C, p_1, \dots, p_n) \in \mathcal{M}_{5,n}$, set once again $D := K - p_1 - p_2$ and consider $C_D \subseteq \mathbb{P}_{\mathbb{C}}^2$. Let

$$V := x_1^5 \mathbb{C}[x_2, x_3]_1 \oplus x_1^4 \mathbb{C}[x_2, x_3]_2 \oplus x_1^3 \mathbb{C}[x_2, x_3]_3 \oplus x_1^2 \mathbb{C}[x_2, x_3]_4 \oplus \langle x_1, x_2 \rangle \mathbb{C}[x_2, x_3]_5$$

and define the incidence variety

$$X' := \{ (f, S_0, \dots, S_4, A_3, \dots, A_n) \in \mathbb{P}(V) \times \mathbb{P}_5^0 \times (\mathbb{P}_{\mathbb{C}}^2)^{n-2} \mid \\ N_i := v(S_i) \text{ is a node on } \{ f = 0 \}, i = 1, \dots, 5, f(A_j) = 0, j = 3, \dots, n \}$$

with projection $\pi: X \rightarrow \mathbb{P}_5^0 \times (\mathbb{P}_{\mathbb{C}}^2)^{n-2}$. Combining the proofs of Lemma 2.5.1 and Lemma 5.3.1 one obtains the following

Lemma 5.5.1. *There exists an open non-empty subset $\mathcal{U} \subseteq (\mathbb{P}_{\mathbb{C}}^2)^{n-2}$ such that the scheme $X := X' \cap (\mathbb{P}(V) \times \mathbb{P}_5^0 \times \mathcal{U})$ is a subbundle of $\mathbb{P}(V) \times \mathbb{P}_5^0 \times \mathcal{U}$ with typical fibre $\mathbb{P}_{\mathbb{C}}^{12-n}$ over $\mathbb{P}_5^0 \times \mathcal{U}$.*

Moreover the general point in X corresponds to an irreducible sextic curve through the points E_1, E_2 , containing exactly five nodes on $\widehat{C}$ and no other singularities. $\square$

As above we conclude $\mathcal{M}_{5,n} \approx X/T$.

Finally we are able to prove the following result.

Main Theorem for $g = 5$. *The moduli space $\mathcal{M}_{5,n}$ is rational for $1 \leq n \leq 12$.*

Proof. In all the cases X is contained in a suitable T -linearized projective bundle over a rational base B contained in the trivial T -equivariant projective bundle $\mathbb{P}(V) \times B$. Clearly there exists a T -invariant unisecant $L \subseteq \mathbb{P}(V) \times B$ (i.e. a divisor intersecting the general fibre in a hyperplane): the scheme $L \cap X$ is then a T -invariant unisecant on X , hence X is T -equivariantly birationally equivalent to a T -equivariant vector bundle $\mathbb{C}^{R(n)} \times B$ where $R(1) = R(2) = 10$ and $R(n) = 12 - n$ for $3 \leq n \leq 12$ as shown in Sections 5.3, 5.4, 5.5.

The action of T on the base B is in all cases almost free, thus $\mathcal{M}_{5,n} \approx X/T \approx \mathbb{C}^{R(n) \times B}/T$ is a vector bundle over the base B/T with $R(n)$ -dimensional fibre due to the results of Section 4 of [Do]. It remains then to prove that the unirational variety B/T is actually rational. To this purpose, we distinguish the cases $n = 1, 2$ and $n \geq 3$.

If $n = 1, 2$ then B is a projective linear representation of the connected solvable group T , hence B/T is rational (see [Mi], [Vi]).

If instead $n \geq 3$ then apply the results of [Do], Section 4, to the projective bundle $\mathbb{P}_5^0 \times (\mathbb{P}_{\mathbb{C}}^2)^{n-2} \rightarrow \mathbb{P}_5^0$. Once again, rationality follows and the proof is over. $\square$

REFERENCES

- [A–S] E. Arbarello, E. Sernesi, *The equation of a plane curve*, Duke Math. J. **46** (1979), 469–485.
- [A–C1] E. Arbarello, M. Cornalba, *Footnotes to a paper of Beniamino Segre*, Math. Ann. **256** (1981), 341–362.
- [A–C2] E. Arbarello, M. Cornalba, *Calculating cohomology groups of moduli spaces of curves via algebraic geometry*, Inst. Hautes Études Sci. Publ. Math. **88** (1998), 97–127.
- [Be] P. Belorousski, *Chow rings of moduli spaces of pointed elliptic curves*, P.H.D. thesis, Chicago, 1998.
- [B–F] G. Bini, C. Fontanari, *Moduli of curves and spin structures via algebraic geometry*, Trans. Amer. Math. Soc. **358** (2006), 3207–3217.
- [Ca] F. Catanese, *On the rationality of certain moduli spaces related to curves of genus 4*, Algebraic geometry (I.V. Dolgachev, ed.), L.N.M., vol. 1008, Springer, 1983, pp. 30–50.
- [C–R] M. Chang, Z. Ran, *Unirationality of the moduli spaces of curves of genus 11, 13 (and 12)*, Invent. Math. **76** (1984), 41–54.
- [Do] I.V. Dolgachev, *Rationality of the fields of invariants*, Algebraic Geometry, Bowdoin 1985 (Spencer J. Bloch, ed.), Proceedings of Symposia in Pure Mathematics, vol. 46, A.M.S., 1987, pp. 3–16.
- [E–H] D. Eisenbud, J. Harris, *The Kodaira dimension of the moduli space of curves of genus ≥ 23* , Invent. Math. **90** (1987), 359–387.
- [Fa1] G. Farkas, *The geometry of the moduli space of curves of genus 23*, Math. Ann. **318** (2000), 43–65.
- [Fa2] G. Farkas, *Syzygies of curves and the effective cone of $\overline{\mathcal{M}}_g$* , Preprint arXiv:math. AG/0503498.
- [Fa3] G. Farkas, *Koszul divisors on moduli spaces of curves*, Preprint arXiv:math. AG/0607475.

- [Fi] E. Fischer, *Die Isomorphie der Invariantenkörpern der endlichen Abelschen Gruppen linearen Transformationen*, Nachr. König. Ges. Wiss. Göttingen (1915), 77–80.
- [G–K–M] A. Gibney, S. Keel, I. Morrison, *Towards the ample cone of $\overline{M}_{g,n}$* , J. Amer. Math. Soc. **15** (2002), 273–294.
- [G–P] T. Graber, R. Pandharipande, *Constructions of nontautological classes on moduli spaces of curves*, Michigan Math. J. **51** (2003), 93–109.
- [Hr] J. Harris, *On the Kodaira dimension of the moduli space of curves, II*, Invent. Math. **75** (1984), 437–466.
- [H–M] J. Harris, D. Mumford, *On the Kodaira dimension of the moduli space of curves*, Invent. Math. **67** (1982), 23–97.
- [Ht] R. Hartshorne, *Algebraic geometry*, Springer, 1977.
- [Ig] J. Igusa, *Arithmetic variety of moduli of genus two*, Ann. of Math. **72** (1960), 612–649.
- [Ka1] P.I. Katsylo, *The rationality of the orbit spaces of irreducible representations of the group SL_2* , Math. USSR Izvestiya **22** (1984), 23–32.
- [Ka2] P.I. Katsylo, *The variety of moduli of curves of genus four is rational*, Soviet Math. Dokl. **34** (1987), 388–389.
- [Ka3] P.I. Katsylo, *Rationality of the variety of moduli of curves of genus 5*, Math. USSR-Sb **72** (1992), 439–445.
- [Ka4] P.I. Katsylo, *Rationality of the moduli variety of curves of genus 3*, Comment. Math. Helvetici **71** (1996), 507–524.
- [Lg] A. Logan, *The Kodaira dimension of moduli spaces of curves with marked points*, Am. J. Math. **125** (2003), 105–138.
- [Lo] E. Looijenga, *Cohomology of $\mathcal{M}_3$ and $\mathcal{M}_3^1$* , Mapping class groups and moduli spaces of Riemann surfaces (Göttingen, 1991/Seattle, WA, 1991) (C.F. Bödigheimer and R.M. Hain, eds.), Contemp. Math., vol. 150, A.M.S., 1993, pp. 205–228.
- [Mi] K. Miyata, *Invariants of certain groups. I*, Nagoya Math. J. **41** (1971), 69–73.
- [P–V] V.L. Popov, E.B. Vinberg, *Invariant theory*, Algebraic Geometry IV (A.N. Parshin and I.R. Shafarevich, eds.), Encyclopedia of Mathematical Sciences, vol. 55, 1991.
- [SB1] N.I. Shepherd-Barron, *The rationality of certain spaces associated to trigonal curves*, Algebraic geometry, Bowdoin 1985 (Spencer J. Bloch, ed.), Proceedings of Symposia in Pure Mathematics, vol. 46, A.M.S., 1987, pp. 165–171.
- [SB2] N.I. Shepherd-Barron, *Invariant theory for S_5 and the rationality of M_6* , Compositio Math. **70** (1989), 13–25.
- [SB3] N.I. Shepherd-Barron, *Rationality of moduli spaces via invariant theory*, Topological methods in algebraic transformation groups (New Brunswick, NJ, 1988) (H. Kraft, T. Petrie and G.W. Schwarz, eds.), Progr. Math., vol. 80, 1989, pp. 153–164.
- [Se] E. Sernesi, *Unirationality of the variety of moduli of curves of genus twelve. (Italian)*, Ann. Scuola Norm. Sup. Pisa Cl. Sci.(4) **8** (1981), 405–439.
- [Ve] A. Verra, *The unirationality of the moduli spaces of curves of genus ≤ 14* , Preprint arXiv:math.AG/0402032.
- [Vi] E.B. Vinberg, *Rationality of the field of invariants of a triangular group*, Mosc. Univ. Math. Bull. **37** (1982), 27–29.

GIANFRANCO CASNATI, DIPARTIMENTO DI MATEMATICA, POLITECNICO DI TORINO, C.SO DUCA DEGLI ABRUZZI 24, 10129 TORINO, ITALY

E-mail address: casnati@calvino.polito.it

C. FONTANARI, DIPARTIMENTO DI MATEMATICA, POLITECNICO DI TORINO, C.SO DUCA DEGLI ABRUZZI 24, 10129 TORINO, ITALY

E-mail address: claudio.fontanari@polito.it